



# SCHEDULE OF RECOMMENDED PREVENTATIVE CARE VISITS AND IMMUNIZATIONS

## "1 Week" (2-3 Days Post Discharge from the Hospital)

Lactation Care is Provided

### 2 Weeks

### 2 Months

Pediarix (DTaP Dose 1 of 5, IPV Dose 1 of 4, HepB Dose 1 of 3) -- Hib (Dose 1 of 4) -- Prevnar 13 (Dose 1 of 4) -- Rotarix (Dose 1 of 2)

### 4 Months

Pediarix (DTaP Dose 2 of 5, IPV Dose 2 of 4, HepB Dose 2 of 3) -- Prevnar 13 (Dose 2 of 4) -- Rotarix (Dose 2 of 2)

### 6 Months

Pediarix (DTaP Dose 3 of 5, IPV Dose 3 of 4, HepB Dose 3 of 3) -- Hib (Dose 3 of 4) -- Prevnar 13 (Dose 3 of 4) -- Flu\*

### 9 Months

### 12 Months

MMR (Dose 1 of 2) -- HepA (Dose 1 of 2) -- Prevnar 13 (Dose 4 of 4)

### 15 Months

DTaP (Dose 4 of 5) -- Varivax (Varicella Dose 1 of 2) -- Hib (Dose 4 of 4)

### 18 Months

### 2 Years

HepA (Dose 2 of 2)

### 30 Months

### 3 Years

### 4 or 5 Years

Kinrix (DTaP Dose 5 of 5, IPV Dose 4 of 4) -- ProQuad (MMR Dose 2 of 2, Varicella Dose 2 of 2)

### 12 Years

HPV (Dose 1 of 2)<sup>^</sup> -- Tdap (Dose 1 of 1) -- Menveo (Dose 1 of 2)

### 16 Years

Menveo (Dose 2 of 2) -- Bexsero (Dose 1 of 2)<sup>^^</sup>

**Pediarix** -- DTaP/IPV/HepB

**DTaP** -- Diphtheria/Tetanus/Acellular Pertussis

**IPV** -- Polio

**Hib** -- Haemophilus Influenzae

**Prevnar 13** -- Pneumococcal

**Rotarix** -- Rotavirus

**MMR** -- Measles, Mumps, and Rubella

**Varivax** -- Varicella (Chicken Pox)

**Kinrix** -- DTaP/IPV

**HPV** -- Human Papillomavirus

**Menveo** -- Meningococcal

**Bexsero** -- Meningitis B

\* -- Flu vaccine may be given starting at 6 months, then annually. If the 1st dose is given before 8 years of age, then a booster is needed at least 1 month later.

<sup>^</sup> -- A second dose of HPV is required 6 months after the first. This can be delivered at a nurse-only visit. If the HPV series is initiated on or after age 15, three doses are required. The second dose should be given 1-2 months after the first dose, and the third dose should be given 6 months after the first dose.

<sup>^^</sup> -- For patients that Men B is recommended for, a second dose is required 1 month after the first. This can be delivered at a nurse-only visit.